

Treatment:

Started CPR

20 seconds later, a normal sinus rhythm returned

Oxygen administered

Pt. had no memory of the event


Had 3 brief periods of emesis

1 hr. later, procedure performed

Next time, pt. pre-medicated the night before and 1 hr. before

Prevention:

- Know the patient (2-4% are really scared)
- Supine position
- Have a monitor available (pulse, BP)
- Administer oxygen
- Pre-medicate patients at risk



DOCS Education courses are either Board-approved or meet Board training requirements for oral conscious sedation in the states and provinces shown in orange.

For more information on regulations in your state please visit SedationRegulations.com or call (888) 316-8327.

Bottom Line with triazolam:

- 0.25 mg or 0.50 mg administered orally or sublingually one hour before the procedure will decrease:
 - anxiety,
 - the perception of pain, and
 - postoperative recall of discomfort

without significant adverse effects. Overdose symptoms develop with four times the maximum therapeutic dose (0.50 mg) or 2 mg.

Case Report: Patient given twice the recommended amount.

Case Report: Patient given twice the recommended amount.

Dear Karl,

It was a pleasure to see you and Niles in Arizona.


I found a sedation course through Baylor University that costs a fraction of the DOCS course. Here is the link (\$695).

<http://www.arcmesa.org/baylor/liveprogram.aspx>

Thanks again for a great course and I look forward to hopefully seeing you again one of these days.

Sincerely,

Milad Bozorgnia



Dental Organization for Conscious Sedation
80 Million North Americans are Too Fearful to Enjoy the Benefits of Complete Dental Care. You can help!

Welcome to a new world of comfort for your patients!

Patients Speak

COURSES

Dental Organization Courses

- "Adult Oral Sedation"
- "N2O-Enteral Conscious Sedation"
- "ANXIOLYSIS - BEYOND VALIUM"
- "The Next Level"
- "Complex Adult Oral Sedation"

Royal Academy Courses

- "Nitrous Oxide"
- "Advanced Sedation Marketing"
- "ACLS"
- "Medical Emerg/In-Office"

www.sedation1.com

LINKS **MEMBERSHIP**

Notes from Dr. Silverman's "DOCS" course, continued:

- Romazicon: 0.2mg (0.1mg/cc) initial dose ½ inch in tissue just off midline under tongue into venous plexus. 2nd dose in 2-3 min. Can cause seizures (if on medications for seizures) or agitation. 10 cc max. Can have re-sedation.
- No smoking or caffeine 12 hrs. before
- Patient should always respond to verbal commands and physical stimuli.

Notes from Dr. Silverman's "DOCS" Course:

- Likes 6 hr. fast before appointment. Avoids erratic or incomplete absorption.
- Need to wake up more? Give some juice.
- Older person needs less medication.
- Contraindications: pregnancy, pt. who premeds with alcohol, erythromycin, Tagamet, antacids, CNS depressants, AIDs patients on protease inhibitors, cyclosporin, grapefruit juice, St. John's Wort, anti-fungal meds patients under 18.

Fred C. Quarnstrom's Home Page: Microsoft Internet Explorer

File Edit View Favorites Tools Help

Address <http://faculty.washington.edu/quarn/>

Fred C. Quarnstrom, D.D.S.

Dental Anesthesiologist
F.A.D.S.A., F.A.G.D., F.L.C.D., C.D.C.
Diplomate, American Dental Board of Anesthesiology
Certified Dental Consultant, American Association of Dental Consultants

*Special procedure not a specialty of the American Dental Association

Office Address:
3051 Breacon Ave. So., Seattle, WA 98144
Telephone: (206) 329-0500 Fax: (206) 329-0538, e-mail quarn@washington.edu

[Beacon Hill Dental Associates](#)
[Dental patient newsletters](#)
[Halcion oral sedation manual](#)
[Risk of nitrous oxide exposure](#)
[Astrenomy](#)

faculty.washington.edu/quarn/

**END
OF
SEDATION**